

Multipurpose Broad Spectrum Farm Disinfectant





effective AT 1% against livestock disease-causing organisms



STABLE AND EFFECTIVE in hot and cold temperatures



EXCELLENT FOAM ADHERENCE on surfaces



NON-CORROSIVE excellent materials compatibility at 1%



READILY
BIODEGRADABLE
at 82% according
to OCDE 301B

Approvals

Approved for disinfecting housing, livestock transport equipment (PT3) and surfaces in contact with foodstuffs and animal feed (PT4).

Composition

- Alkyl dimethyl benzyl ammonium chloride (ADBAC) 32.75%
- Glutaraldehyde 10%
- Excipients: surfactants, sq.

Product Presentations

1L, 5L, 10L, 25L, 60L and 200L

Physical and Chemical Properties


- Colour: amber solution which may intensify over time.
- Fragrance: lemon-scented
- Density ≈ 1.024
- pH of the concentrate ≈ 2.5
- pH of 1% solution ≈ 3.8

Application & Use Guidance

| Disinfection Task | Dilution Rate | Application |
|---------------------------|--|---|
| Surfaces and Equipment | 1% or 1:100 (10 ml of TH5® concentrate per 1 litre of water) | Using a pressure washer or other mechanical sprayer, apply at a rate of 300 ml/m² (or to point of "run-off") using either a foam or spray lance. Leave the surface to dry. Surfaces in contact with foodstuffs or animal feed should be rinsed with clean water. Wait until the building is completely dry before placing animals (24 to 48 hours). |
| Vehicles | | Working from the top down, apply TH5® solution using a pressure washer or other mechanical sprayer paying particular attention to the wheels, wheel arches, mudguards and any exposed chassis of the vehicle. |
| Boot & Wheel Dips | | Fill the boot or wheel dip with the required volume of TH5® concentrate and water and stir thoroughly. Ensure all persons step into the boot dip and vehicles travel through the wheel dip before entry to site or buildings. Renew disinfectant solution every 2 days or before if heavily soiled. |
| Cold Fogging | 2.5 ml/m ³ | Dilute 1:1 (1 litre of TH5® concentrate with 1 litre of water). Use a mechanical mister to apply the TH5® solution. Leave the building closed for a minimum of 30 minutes and allow mist to fully disperse before re-entering. |
| Thermal Fogging | 1 ml/m ³ | Dilute 1:1 (1 litre of TH5® concentrate with 1 litre of water). Fill the thermal fogging machine with the required volume of TH5® solution. Direct the nozzle of the fogger into the building, with the air intake situated outside of the building, and activate. Leave the building closed for a minimum of 1 hour and allow mist to fully disperse before re-entering. |

Always use Personal Protective Equipment (PPE) when handling and applying chemical concentrates and in-use solutions:

Overall - chemical risk type 4 single use, Nitrile Gloves EN374, Half mask - with canisters ABEK P, Protective glasses or complete mask.



| Number of pigs | Surface area | Volume of TH5®1% solution | Volume of TH5® concentrate |
|-------------------|--------------------|---------------------------------|----------------------------------|
| Per 10 Sows | 240 m ² | 72 litres | 720 ml |
| Per 100 Weaners | 150 m ² | 45 litres | 450 ml |
| Per 100 Fatteners | 300 m ² | 90 litres | 900 ml |



| Floor Type | Surface area | Volume of TH5®1% solution | Volume of TH5® concentrate |
|------------|---------------------|---------------------------------|----------------------------------|
| Dirt | 1500 m ² | 450 litres | 4.5 litres |
| Concrete | 2500 m ² | 750 litres | 7.5 litres |

Thermal Fogging Disinfection Usage Chart

| Floor area: 1000 m ² | House volume | Volume of Water | Volume of TH5® concentrate |
|------------------------------------|---------------------|--------------------|----------------------------------|
| Dirt and concrete floor | 3500 m ³ | 3.5 litres | 3.5 litres |

Bactericidal Efficacy data

| Bacteria | Test Method | Dilution Rate (%) |
|---------------------------------|--|----------------------|
| Campylobacter jejuni | EN 1656 (4 °C & 10 °C – 5 mins) | 0.1 |
| Enterococcus hirae | EN 14349 (4 °C- 30 mins) | 0.25 |
| Escherichia coli | EN 1656 (4 °C & 10 °C – 30 mins) | 0.1 |
| Mycoplasma hyopneumoniae | NFT 72-180 (4 °C & 10 °C – 30 mins) | 1 |
| Ornithobacterium rhinotracheale | EN 1656 (10 °C - 5 mins) | 0.1 |
| Pseudomonas aeruginosa | EN 14349 (10 °C - 30 mins) | 0.15 |
| Salmonella Enteritidis | EN 1656 (4 °C - 60 mins) | 0.1 |
| | UK DEFRA (4 °C - 30 mins) | 2.85 |
| Salmonella Gallinarum | EN 1656 (10 °C - 30 mins) | 0.2 |
| Salmonella Typhymurium | EN 1656 (4 °C - 30 mins) | 0.1 |
| Salmonella Virchow | EN 1656 (10 °C - 30 mins) | 0.1 |
| Streptococcus suis | EN 1656 (10 °C - 30 mins) | 0.05 |

Fungicidal & Yeasticidal Efficacy Data

| Fungi & Yeast | Test Method | Dilution Rate (%) |
|-----------------------------|--------------------------------|----------------------|
| Aspergillus brasiliensis | EN 1657 (10 °C - 30 mins) | 0.5 |
| Aspergilus fumigatus | EN 1657 (10 °C - 30 mins) | 1 |
| Candida albicans | EN 16438 (20 °C - 120 mins) | 0.1 |
| Trichophyton mentagrophytes | EN 16438 (10 °C - 60 mins) | 0.5 |
| Trichophyton verucosum | EN 16438 (10 °C - 60 mins) | 1 |

Virucidal Efficacy data

| Disease/Related Condition | Test Method | Dilution Rate (%) |
|---|------------------------------------|----------------------|
| African Swine Fever (ASF) | EN 14675 (10 °C - 30 mins) | 0.3 |
| Aujeszky's disease | NFT 72-180 (10 °C - 30 mins) | 0.1 |
| Avian Reovirus | EN 14675 (10 °C - 30 mins) | 0.2 |
| Classical Swine Fever (CSF) | EN 14675 (4 °C - 30 mins) | 0.05 |
| Feline Coronavirus | EN 17122 (4 °C - 60 mins) | 0.5 |
| Foot and Mouth Disease (FMD) | UK DEFRA (4 °C - 30 mins) | 0.5 |
| Infectious Bronchitis (IB) | NFT 72-180 (20 °C - 30 mins) | 0.2 |
| Infectious Bursal Disease (IBD) | EN 14675 (10 °C - 30 mins) | 0.5 |
| Influenza virus: H1N1, H7N1, H7N9, H5N1 | EN 14675 (10 °C - 30 mins) | 0.1 |
| Marek's Disease | NFT 72-180 (10 °C - 30 mins) | 0.1 |
| Newcastle Disease (ND) | EN 14675 (4°C - 30 mins) | 0.25 |
| | UK DEFRA (4°C - 30 mins) | 1.33 |
| Porcine Circovirus type 2 (PCV2) | EPA Compliant (20 °C - 30 mins) | 1 |
| Porcine Epidemic Diarrhoea (PED) | EPA Compliant (10 °C - 60 mins) | 0.5 |
| Porcine Parvovirus | EN 17122 (10 °C - 120 mins) | 1 |
| Porcine Reproductive and Respiratory Syndrome (PRRS) | EN 14675 (10 °C - 30 mins) | 0.1 |
| Poxvirus | NFT 72-180 (20 °C - 30 mins) | 0.1 |

LANXESS

THESEO France 200 avenue de Mayenne 53000 Laval – France

Tel: +33 (0)2 43 67 96 96 Fax: +33 (0)2 43 67 97 00 contact.LBSFrance@lanxess.com biosecurity@lanxess.com biosecuritysolutions.lanxess.com lanxess.com This information and our technical advice – whether verbal, in writing or by way of trials – is subject to change without notice and given in good faith but without warranty or guarantee, express or implied, and this also applies where proprietary rights of third parties are involved. Our advice does not release you from the obligation to verify the information currently provided - especially that contained in our safety data and technical information sheets - and to test our products as to their suitability for the intended processes and uses. The application, use and processing of our products and the products manufactured by you on the basis of our technical advice are beyond our control and, therefore, entirely your own responsibility. Our products are sold in accordance with the current version of our General Conditions of Sale and Delivery.

The specified uses and registered claims for the product may vary from country to country. Please contact LANXESS to verify country-specific approved uses.

Use biocides safely. Always read the label and product information before use. Dangerous product. Respect precautions for use. For professional use only.

©2023 LANXESS. TH5®, LANXESS, the LANXESS Logo and any associated logos are trademarks or copyrights of LANXESS Deutschland GmbH or its affiliates. All trademarks are registered in many countries worldwide.